

**(carcinoma of the bronchus, bronchogenic carcinoma)****DEFINITION**

1. Cancer is a term which embraces a large number of different diseases, the common feature of which is a malignant tumour. This is a growth (neoplasm) which is not circumscribed but which infiltrates the surrounding tissues and metastasises (spreads to other sites in the body), thereby producing secondary deposits. Any tissue in the body may be affected.
2. Cancers are classified according to the tissue of origin. **Carcinoma** arises from epithelial tissue and sarcoma from connective tissue. The suffix **blastoma** implies a tumour of embryonic origin.
3. Cancer of the lung is a malignant tumour of the lung. The World Health Organisation (1981) classification lists the following four main tumour types (incidence given in brackets):
  - 3.1. **epidermoid or squamous cell carcinoma (35%)**
  - 3.2. **small or oat cell carcinoma (24%)**
  - 3.3. **adenocarcinoma (21%)**
  - 3.4. **large cell carcinoma (15%)**

The remaining malignant lung tumours include mesothelioma, carcinoids and muco-epidermoid carcinoma.

4. The site of the tumour is variable. There appears to be a predilection for the bifurcations of the bronchi. A significant number of lung tumours arise in the periphery of the lung, where there is a preponderance of adenocarcinomas and large cell tumours.
5. Squamous cell tumours are relatively slow growing and have a low incidence of haematogenous spread. In contrast, small cell tumours are fast growing and metastasise early. Adenocarcinomas and large cell tumours behave in an intermediate fashion.

**CLINICAL MANIFESTATIONS**

6. In about 5% of cases of bronchial cancer the patient is symptom-free, the disease being detected only by routine chest X-ray.
7. Clinical features may be due to tumour growth within the lung, invasion of tumour to structures outwith the lung but within the thorax, or lymphatic or haematogenous metastatic spread. Endocrine, metabolic or neurological syndromes may be present.

8. The most usual presenting symptom is cough, often said to be brassy. Sputum expectoration is common, the sputum usually being viscid or purulent. 5% of cases present with haemoptysis, and in about half the cases haemoptysis occurs at some stage in the disease. Wheeze, stridor and dyspnoea may be present. Vague chest discomfort is common. There may be pleuritic or bone pain. Arm and shoulder pain occurs in tumours of the superior pulmonary sulcus. Characteristic non-specific symptoms include lack of energy, loss of interest and pyrexia.
9. Direct spread of the tumour to intrathoracic structures may cause well recognised syndromes. Invasion of the left recurrent laryngeal nerve produces vocal cord paresis. Horner's syndrome with miosis, anhidrosis and enophthalmos results from involvement of the last cervical and first thoracic segments of the sympathetic trunk. Compression of the superior vena cava produces suffusion of the head, neck and upper trunk.
10. Other possible features are dysphagia, arrhythmias, cardiac failure from heart and pericardial metastases, and paradoxical movement of the diaphragm from phrenic nerve invasion. Infiltration of the ribs and pleura, and spontaneous pneumothorax, may occur. There may be direct spread to the other lung.
11. Metastatic lesions may occur in any site. Lymphatic spread often involves the scalene node. Bony metastases occur particularly in the ribs, femora, vertebrae and humerus. 4% of patients present with brain secondaries, and these are clinically apparent at some time in the course of the disease in about 1/3 of cases. Spread to the liver, adrenals, kidneys and skin also occurs.
12. Some bronchial cancers produce ectopic hormones. These result in a number of syndromes whose clinical features reflect the hormone produced. Such ectopic hormones include ADH, ACTH, parathormone, thyroid and melanocyte-stimulating hormone.
13. Abnormalities of the central, peripheral and autonomic nervous systems and the muscles may be found, resulting in neuropathies and myopathies. Specific patterns are recognised, and these features often regress on treatment of the parent tumour.
14. Finger clubbing and the related hypertrophic pulmonary osteoarthropathy may occur. These may precede diagnosis of the tumour.
15. Normochromic normocytic anaemia is usual. Polycythaemia can occur and this leads to hypercoagulability states, sometimes resulting in thrombophlebitis.

### **Characteristics of neoplasia**

16. Tumours arise when tissue growth becomes insensitive to normal control mechanisms. Usually, cells multiply in response to a reduction in tissue cell density, the process ceasing when normal tissue architecture is restored. Tumours arise from a common ancestral cell which – years before clinical disease becomes apparent – begins to reproduce inappropriately. Malignant transformation results from mutation in the cell's genetic material.

17. These genetic mutations change the amount or the activity of the protein product. Two particular gene classes play major roles in triggering cancer. Normally **proto-oncogenes** encourage cell growth and reproduction: **tumour suppressor genes** oppose it. Mutated proto-oncogenes become **carcinogenic oncogenes**, driving excess cell multiplication. Inactivated tumour suppressor genes contribute to cancer by depriving the cell of its braking system. For cancer to develop, mutations need to occur in several of the original cell's growth control genes.
18. Cells insensitive to growth control generate, within tissues, subpopulations of cells whose subsequent progress depends on selection pressure, favouring the emergence of cells with the greatest growth potential. It is this process which determines the histology of a tumour and its clinical course. Tumour cells usually retain the characteristics of their tissue of origin. However, tissue differentiation involves the synthesis of molecules not essential to cell survival, and so loss of the parent tissue markers may confer growth advantage. Dedifferentiation in a tumour implies high growth rate and high malignancy. Tumours can also metastasise (migrate via the lymph or blood stream) to other parts of the body, producing metastatic deposits.

## **AETIOLOGY**

### **General risk factors for all cancers**

19. Much is now understood about the origins and mechanisms of cancer in general, but the precise causes of individual tumours often remain unknown.

### **Age**

20. Some risk of cancer occurs at every age, but the risk for particular types varies at different ages. Most commonly there is a progressive increase in incidence from childhood to old age, (cancers of the skin, lung, gastrointestinal and genitourinary tracts, multiple myeloma and chronic lymphatic leukaemia). The rate of increase is typically proportional to the 4<sup>th</sup> or 5<sup>th</sup> power of age. Less commonly there may be a peak in early life, with decline to zero thereafter, eg retinoblastoma or neuroblastoma.

### **Gender**

21. The incidence of cancer in relation to gender is not straightforward and it is changing – probably due to lifestyle and occupational changes. At the beginning of the last century, overall cancer was more common in women due to the frequency of cancer of the cervix and the rarity, in both men and women, of cancer of the lung. Now, in developed countries, cancer is more common in men.

## Latent period

22. There is a delay between initial exposure to carcinogen and clinical disease. With short intense exposures, eg to ionising radiation at Hiroshima, solid tumours increase in incidence for 15-20 years. Incidence may then continue to rise, level off or decline, depending on the tumour type. For acute leukaemia there is a peak incidence at about 5 years, and very few cases appear after 30 years. Short intense exposure to carcinogens is exceptional and, where exposure is chronic and prolonged, eg due to occupation, smoking or sunlight, cancer incidence increases with exposure duration. Precisely how this operates and how subsequent risk is affected by removal from the exposure is unknown.
23. Clinical cancer is the end result of a multistage process involving initiating and promoting agents. If the carcinogen is an initiating agent, eg asbestos, rather than a substance influencing a later stage nearer clinical manifestation, eg cigarette smoking, cancer incidence in the population may continue to rise, albeit more slowly, for a considerable time after exposure to the carcinogen has ceased.

### **Risk factors in the individual case of all cancers**

24. The main factors that determine whether a particular individual develops cancer relate to constitution and exposure to environmental factors.

### **Genetics**

25. The close connection between certain chromosomal abnormalities associated with recognised clinical syndromes and subsequent tumour development, eg polyposis coli and cancer of the large bowel, and xeroderma pigmentosum and skin tumours, confirms that an individual's genetic make-up has an effect on his susceptibility to cancer.
26. Many studies have looked at cancer rate in the families of individuals with the disease. There appears to be no material tendency for cancer in general to cluster in families and no genes have been identified that increase the risk of cancer in all tissues. However, all common cancers do cluster in families to some extent – the risk of a sibling of a patient developing a tumour at the same site is twice normal. This might be due to genetic susceptibility, but could equally well reflect lifestyle, eg diet, hygiene or a common legacy of infections in early life.

### **Environmental factors**

27. Our knowledge of the environmental causes of cancer relies on animal laboratory investigation and human epidemiology, with the two approaches complementing each other. Since there are features common to most cancers, there are factors which can cause cancer at all or many sites. Present evidence confirms the importance of life-style factors in cancer causation.

## **Tobacco smoke**

28. Cigarette smoking is thought to cause 30% of all cancer deaths and has been conclusively linked to cancer of the lung, upper respiratory tract, oesophagus, bladder, stomach, liver, kidney, chronic myeloid leukaemia. It may also cause cancer of the colon and the rectum. Relevant factors include the number of cigarettes smoked, tar content, age at smoking onset and duration of habit.

## **Diet**

29. There is good evidence that some common cancers would be less common if diet were modified. Animal fat consumption, particularly red meat, high salt intake and ingestion of very hot beverages and food have all been linked to specific cancers. Similarly, what is **not** in the diet may be important. Low consumption of vegetables and fruit, in the presence of high calorie intake, is associated with several different tumour types, eg childhood obesity and cancer of the breast and prostate, adult obesity and endometrial cancer. Consumption of alcohol (particularly along with cigarettes) increases the risk of cancer of the upper respiratory and digestive tracts. There is evidence that as little as two drinks a day may contribute to breast, colonic and rectal cancer. In total, diet is considered to account for 30% of all cancer mortality in developed countries, alcohol for a further 3% and salt for 1%.

## **Radiation**

30. Radiation is difficult to avoid and, in total, radiation of all types causes 2% of all cancer deaths. Most of these deaths result from natural sources, particularly sunlight, UVB.
- **UVB radiation** causes 90% of all skin cancers, including basal cell cancers, malignant melanoma and squamous cell carcinoma.
  - **Electromagnetic radiation** as a cause of cancer has been the subject of several recent studies. The results are confusing and inconsistent and reported associations may not be causal. It is of two main types:
    - i. **Extremely low frequency fields**, eg power lines and household appliances. Basic science confirms that these radiations are of too low frequency to initiate cancer causing genetic mutation as they are of insufficient energy to ionise molecules.
    - ii. **Radiofrequency electromagnetic radiation**, eg cellular telephones, microwaves and living creatures. Although more energetic than i. (above), they are still unable to cause molecular ionisation.

In conclusion, at this date there is no good scientific evidence that electromagnetic radiation causes cancer. Any possible association remains hypothesis.

- **Ionising radiation**

Ionising radiation can penetrate animal tissues and damage DNA and theoretically has the power to produce cancer in most tissues. The actual risk due to exposure to ionising radiation may, however, be different. It is often overestimated and not evidence-based. Amongst Japanese residents of Hiroshima and Nagasaki who survived more than a year after detonation, only 1% have died of tumours.

Studies of humans exposed to high dosage of ionising radiation, eg the Japanese atomic bomb survivors or individuals medically irradiated for tumours, have shown an increased incidence of cancer due to that exposure. There is, however, no firm evidence from human low-dose epidemiological studies which unequivocally demonstrates an increase in cancer incidence. This may be due to the very large size of study population which would be needed to demonstrate an increased incidence.

For radiation protection purposes it is, therefore, accepted that there is no threshold level below which no carcinogenic effect is produced and the risk of a cancer developing is extrapolated on a dose-proportional basis from high to low doses and dose rates.

All humans are constantly exposed to ionising radiation from both the natural environment and man-made products. The natural sources include cosmic radiation from space, radiation from the ground, and from inhaled and ingested materials. Air travel and mining both increase exposure to background radiation. Radiation originating in the body comes mainly from potassium, while lungs are exposed through radon in inhaled air. Man-made radiation comes from medical uses, past atomic tests, man-made products and radioactive waste.

Natural radiation differs depending on location. In the UK the average annual dose is less than 2,000 microsieverts. There is, however, a considerable range; it may rise to 8,000 microsieverts in some areas and to 100,000 in some homes. The UK average annual dose from man-made sources in total is less than 300 microsieverts and, again, there may be variation.

From 1952 to 1958 the UK carried out 21 atmospheric nuclear tests in the Pacific Ocean. The locations were chosen because of their isolation and low natural radiation level. On average the Christmas Island annual background radiation is less than 700 microsieverts.

### **Radiation dose**

31. The effects of ionising radiation depend on the exposure size of the accumulated dose. A discussion of radiation dose is at Annex A.

## Therapeutic drugs

32. About 20 agents, not all of which are in current use, are known to cause cancer. Potential carcinogens may still be used if the hazard is judged to be less than the chance of saving a life, eg certain cancer drugs. Close scrutiny is kept on drug hazards and the position of oestrogens in hormone replacement therapy (HRT), known to cause endometrial cancer, and of the oral contraceptive pills, which have been associated with carcinoma of the cervix, breast and hepatoma, is closely monitored. Together, prescribed drugs are held responsible for less than 1% of all fatal cancers.

## Occupation

33. Historically, study of occupational exposures has identified many important carcinogens. Material or process modification and, latterly, health and safety statute have removed many potential hazards in the developed world. However, the long latent period of cancer means that a considerable time will be required for the effects of industrial carcinogens to be eliminated and, equally, that new hazards may remain unsuspected for a long time. At present overall, occupation is considered responsible for 2-3% of all fatal cancers in developed countries. Particularly important occupational carcinogens are asbestos dust exposure, exposure to combustion products of fossil fuels and ionising radiation.

## Pollution

34. Investigation of the relation between environmental pollution – air, soil and water – and cancer is difficult because of the widespread nature of pollution and similar risk to people over a wide geographical area. It is generally accepted that, in the UK at the beginning of the last century, air pollution via combustion may have contributed to a few percent of lung cancers. Over the last 30 years, with increasing statute on pollution reduction, this has become much less common. Advances in chemical analysis have allowed recent interest in pollution of soil and water as possible cancer risks.
35. Another complicating factor in accurately attributing risk of cancer to individual external agents is **interaction**. Some carcinogenic agents act together to produce effects much greater than the sum of the separate individual effects, eg smoking and asbestos in relation to cancer of the lung, smoking and alcohol in relation to carcinoma of the oesophagus, and aflatoxin and hepatitis B infection in cancer of the liver.

## SPECIFIC RISK FACTORS FOR CANCER OF THE LUNG

36. Although rare until the 1920s, in England and Wales recent cancer registry statistics show that cancer of the lung is now responsible for 16.8% of all cancers and 24% of cancer deaths. The sex ratio male/female is 3.5:1. In developed countries lung cancer is the most important cancer of men, and the incidence in women is rising.

37. Smoking of any tobacco product increases the risk of lung cancer. In the early 1990s smoking caused 90% of all lung cancers in UK. This risk is greatest for cigarette users, and the risk increases with the number of cigarettes smoked and the earlier the age at which smoking began. If individuals stop smoking, the risk declines and, by 15 years, is 11% of that of continuing smokers. Squamous carcinoma is most closely linked to smoking and adenocarcinoma least. Oat cell cancers occupy an intermediate position.
38. There are recognised occupational causes of lung cancer. Working with asbestos may cause lung cancer. Due to interaction the risk is much increased if the worker also smokes cigarettes. The link with asbestos is less specific than that for mesothelioma, and clinical or radiological evidence of asbestos exposure in the form of pleural plaques, bilateral diffuse pleural thickening or asbestosis will be present. The mining of pitch-blende and uranium, haematin and fluorspar, working in foundries, gas and coke ovens (where polycyclic aromatic hydrocarbons are thought to be involved), and the nickel, chromate, arsenical salt and mustard gas manufacturing industries, are all associated with the development of lung cancer.
39. No causal link has been confirmed in the last 30 years between atmospheric pollution and the incidence of lung cancer. Studies into this are very difficult to control as they invariably include multiple interdependent possible causal factors, eg smoking.
40. Lung cancer is strongly linked to exposure to ionising radiation. If this is the cause, it is most usually of the oat cell type and carries a poor prognosis.
41. Peripheral tumours may arise in residual scar tissue from infection or inflammation.
42. Cancer of the lung has not been shown to be caused by climatic extremes, trauma, physical or mental stress, or lowered resistance arising from hardship or other diseases. There is no evidence that its progress is dependent on external factors, other than medical treatment.

## **CONCLUSION**

43. Cancer of the lung is a malignant tumour in which a number of aetiological factors may be present, including cigarette smoking and certain industrial processes. The course of the condition is unaffected by environmental factors other than those involved in its treatment.

## **REFERENCES**

Boice J D. Studies of Atomic Bomb Survivors. JAMA 1990;264:622-623.

Darby, S C, Kendall G M, Fell T P et al. A summary of mortality and incidence of cancer in men from the United Kingdom who participated in the United Kingdom's atmospheric nuclear weapon tests and experimental programmes. BMJ 1990;296:332-338.

Doll R and Peto R. Epidemiology of Cancer. In: Eds. Weatherall D J, Ledingham J G G and Warrell D A. Oxford Textbook of Medicine. 3<sup>rd</sup> Ed. 1996. Oxford. Oxford University Press. 4.95-4.123.

Doll R. Epidemiology of Human Neoplasia. In: Eds. McGee J O'D, Isaacson P G and Wright N A. Oxford Textbook of Pathology. 1992. Oxford. Oxford University Press. 679-694.

Harnden D G, Lorenzen J, Pusztai L and McGee J O'D. Carcinogenesis. In: Eds. McGee J O'D, Isaacson P G and Wright N A. Oxford Textbook of Pathology. 1992. Oxford. Oxford University Press. 633-678.

Hausen H, et al. Tumours of the Trachea and Lung. In Eds: Peckham M, Pinedo, H and Veronesi U. Oxford Textbook of Oncology. 1<sup>st</sup> Ed. 1995;11.1:1533-1557.

Pierce DA et al. Studies of the mortality of atomic bomb survivors. Report 12, part 1. Cancer: 1950-1990. Rad. Res. 1996;146:1-27.

Schimizu Y, Schull W J and Kato H. Cancer risk among Atomic Bomb Survivors: the RERF Life Span Study. JAMA 1990;264:601-604.

Spiro S G. Tumours of the lung a. lung cancer. In: Eds Weatherall D J, Ledingham J G G and Warrell D A. Oxford Textbook of Medicine. Oxford: Oxford University Press. 3<sup>rd</sup> Ed. 1976;17.13.1:2879-2893.

Thompson DE et al. Cancer incidence in atomic bomb survivors Part II: Solid Tumours, 1958-1987. Rad. Res. 1994;137:S17-67.

March 2000

## Radiation dose

1. The first definition of a unit of radiation dose was made in 1928 by the International Congress of Radiology. The roentgen (R) was defined as that quantity of radiation which produces in 1 cm of air one unit of charge of either sign, thus defining a unit of exposure. Units of **absorbed dose**, the actual energy absorbed in the tissue being irradiated are now used. The radiation absorbed dose or **rad** is now cited in SI (Système Internationale) units – joules per kg – of absorbing material. The fundamental unit, 1 joule/kg, is 1 gray (1 Gy), equivalent to 100 rads (R).
2. Different radiation types have greater or lesser effect per unit dose, so they are all expressed relative to the effects of X-rays, ie. a unit equivalent dose is used. To calculate the roentgen equivalent in man (**rem**), the absorbed radiation dose is multiplied by a radiation weighting factor, dependent on type and energy of the radiation. The current SI unit of equivalent dose is the **Sievert**. For X-rays and gamma rays the equivalent dose in sieverts and the absorbed radiation dose in grays are the same. The relationship between the different dose units is:-

1 gray (Gy) = 1 joule/kg = 100 rads (R) = 100 rems (r) = 1 sievert (Sv) = 1,000 millisieverts (mSv) = 1,000,000 microsieverts (microSv). Typical doses of radiation include:

Chest X-ray – 0.02 mSv

Brain scan – 7 mSv

Bone scan – 4 mSv

Average annual UK dose from cosmic rays – 0.26 mSv

Average annual UK dose from gamma rays – 0.35 mSv

Average annual UK dose from natural background radiation – 2.2 mSv

### 3. Effects of total body irradiation

Equivalent dose (Sv)	Effect
Sub lethal to man 0.0001 (0.1 mSv)	Around 2 weeks' natural background radiation, no detectable effect
0.001 (1 mSv)	Around 6 months' natural background radiation, no detectable effect
0.01 (10 mSv)	No detectable effect
0.1 (100 mSv)	Minimal decrease in peripheral lymphocyte count, no clinical effect
1 (1000 mSv)	Mild acute radiation sickness in some individuals (nausea, possible vomiting), no acute deaths, early decrease in peripheral lymphocyte count, decrease in all WBC and platelets at 2-3 weeks, increase in late risk of leukaemia, solid tumours

<b>Equivalent dose (Sv)</b>	<b>Effect</b>
Lethal to man 10 (10,000 mSv)	Severe acute radiation sickness, severe vomiting, diarrhoea, death within 30 days of all exposed individuals. Severe depression of blood cell and platelet production, damage to gastrointestinal mucosa.
100 (100,000 mSv)	Immediate severe vomiting, disorientation, coma, death within hours
1000 (1,000,000 mSv)	Death of some micro-organisms, some insects within hours
10,000 (10,000,000 mSv)	Death of most bacteria, some viruses
100,000 (100,000,000 mSv)	Death of all living organisms, denaturation of proteins

### **Radiation dose limits**

4. Since the days of Marie Curie it has been appreciated that ionising radiation exposure may be hazardous to health. Radiation dose limits were first recommended for ionising radiation exposure in 1928. The statutory limit on the amount of radiation to which the general public may be exposed in excess of natural background radiation and excluding medical exposure is set, from 1 January 2000, at 1 mSv per annum.
5. The most important source of man-made exposure is medical investigation which accounts for 90% of man-made exposure. Average natural background radiation is raised to 2.6 mSv by all man-made exposure. UK estimated exposure, excluding medical investigation, is 0.04 mSv. Other statutory limits include occupational dose limits. From 1 January 2000, these are 20 mSv per annum for classified workers and 6 mSv per annum for unclassified workers.

January 2000